

# Circulation Research

JOURNAL OF THE AMERICAN HEART ASSOCIATION



## Direct Cardiac Reprogramming: From Developmental Biology to Cardiac Regeneration Li Qian and Deepak Srivastava

*Circ Res.* 2013;113:915-921

doi: 10.1161/CIRCRESAHA.112.300625

*Circulation Research* is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2013 American Heart Association, Inc. All rights reserved.

Print ISSN: 0009-7330. Online ISSN: 1524-4571

The online version of this article, along with updated information and services, is located on the  
World Wide Web at:

<http://circres.ahajournals.org/content/113/7/915>

**Permissions:** Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation Research* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

**Reprints:** Information about reprints can be found online at:  
<http://www.lww.com/reprints>

**Subscriptions:** Information about subscribing to *Circulation Research* is online at:  
<http://circres.ahajournals.org/subscriptions/>

This article is in a thematic series on **Recent Advances in iPSC Cell Research**, which includes the following articles:

Steps Toward Safe Cell Therapy Using Induced Pluripotent Stem Cells [*Circ Res.* 2013;112:523–533]

iPSCs in Cardiovascular Drug Discovery [*Circ Res.* 2013;112:534–548]

Immunogenicity of Pluripotent Stem Cells and Their Derivatives [*Circ Res.* 2013;112:549–561]

Progress in the Reprogramming of Somatic Cells [*Circ Res.* 2013;112:562–574]

**Direct Cardiac Reprogramming: From Developmental Biology to Cardiac Regeneration**

Use of iPSCs to Generate Cardiac Myocytes and Repair Infarcted Myocardium

Genomic Stability Issues Associated With the Reprogramming Process

Differentiation of iPSCs Into Mature Myocytes

*Shinya Yamanaka, Guest Editor*

## Direct Cardiac Reprogramming From Developmental Biology to Cardiac Regeneration

Li Qian, Deepak Srivastava

**Abstract:** Heart disease affects millions worldwide and is a progressive condition involving loss of cardiomyocytes. The human heart has limited endogenous regenerative capacity and is thus an important target for novel regenerative medicine approaches. Although cell-based regenerative therapies hold promise, cellular reprogramming of endogenous cardiac fibroblasts, which represent more than half of the cells in the mammalian heart, may be an attractive alternative strategy for regenerating cardiac muscle. Recent advances leveraging years of developmental biology point to the feasibility of generating de novo cardiomyocyte-like cells from terminally differentiated nonmyocytes in the heart in situ after ischemic damage. Here, we review the progress in cardiac reprogramming methods and consider the opportunities and challenges that lie ahead in refining this technology for regenerative medicine. (*Circ Res.* 2013;113:915-921.)

**Key Words:** fibroblasts ■ heart diseases ■ myocytes, cardiac ■ reprogramming

Cardiomyocyte death is associated with diverse forms of heart disease, which is the leading cause of morbidity and mortality in industrialized countries. Myocardial infarctions are a major public health problem and affect >1 million Americans yearly. A tremendous loss of cardiomyocytes through apoptosis and necrosis underlies both acute and chronic myocardial infarction. To compensate for the death of cardiomyocytes in the injured area, scar tissue is formed by activated fibroblasts. In addition, heart diseases related to pressure overload, such as hypertension or valvular disease, involve the death of cardiac muscle cells, but the progression is slow and usually occurs during a long period of time.

Human aging is associated with an irreversible loss of cardiomyocytes, which may account for the increased vulnerability of aged hearts to various risk factors.<sup>1</sup>

Unlike amphibian and fish hearts,<sup>2</sup> the human heart has limited regenerative potential. Although postnatal vertebrate cardiomyocytes undergo some degree of cell renewal, as suggested by transgenic mouse experiments<sup>3–5</sup> and a more recent human radiocarbon dating study,<sup>6</sup> the rate of cardiomyocyte renewal is low. A genetic fate map study involving mice indicated that the endogenous regenerative capacity of the adult heart, albeit limited, largely comes from the differentiation of cardiac progenitor cells rather than replacement by existing cardiomyocytes

Original received December 6, 2012; revision received January 15, 2013; accepted February 4, 2013. In July 2013, the average time from submission to first decision for all original research papers submitted to *Circulation Research* was 13.24 days.

From the McAllister Heart Institute, Department of Pathology and Laboratory Medicine, School of Medicine, University of North Carolina at Chapel Hill, Chapel Hill (L.Q.); Gladstone Institute of Cardiovascular Disease and Roddenberry Center for Stem Cell Biology and Medicine at Gladstone, San Francisco, CA (D.S.); Department of Pediatrics and Biochemistry and Biophysics, University of California, San Francisco, CA (D.S.).

Correspondence to Deepak Srivastava, MD, Gladstone Institute of Cardiovascular Disease, 1650 Owens St, San Francisco, CA 94158. E-mail dsrivastava@gladstone.ucsf.edu

© 2013 American Heart Association, Inc.

*Circulation Research* is available at <http://circres.ahajournals.org>

DOI: 10.1161/CIRCRESAHA.112.300625


### Nonstandard Abbreviations and Acronyms

<b>GMT</b>	Gata4, Mef2c, and Tbx5
<b>iCM</b>	induced cardiomyocyte
<b>iPSC</b>	induced pluripotent stem cells
<b>miRNA</b>	microRNA

through cell division.<sup>7,8</sup> A major challenge moving forward is to identify such cardiac progenitor pools in the adult heart and promote their expansion and differentiation potential in vivo. In contrast, Porrello et al<sup>9</sup> reported the neonatal mouse heart has a remarkable regenerative capacity within the first 7 days of life, similar to that of the zebrafish heart. Genetic fate mapping suggests that myocyte proliferation is the main mechanism for this regeneration,<sup>9</sup> which is reminiscent of what has been found in the adult zebrafish heart.<sup>10,11</sup> Furthermore, in a recent report using stable isotope labeling with genetic fate mapping, in combination with multi-isotope imaging mass spectrometry, the authors concluded that pre-existing cardiomyocytes are the dominant source of myocyte renewal in the adult mouse heart with or without injury.<sup>5</sup> Cardiomyocyte proliferation has been reported to be enhanced by overexpressing cyclin D2<sup>12,13</sup> or administering factors, such as periostin,<sup>14</sup> fibroblast growth factor-1,<sup>15</sup> or neuregulin 1,<sup>16</sup> in mouse models of myocardial infarction. Most recently, addition of microRNA-199a and microRNA-590 also stimulated cardiomyocyte proliferation in vitro and in vivo.<sup>17</sup> Thus, there is increasing evidence that adult mammalian cardiomyocytes have the capacity to proliferate, but whether approaches to enhance this feature are sufficient to compensate for the functional loss of damaged myocardium awaits confirmation.

Another strategy to repair an injured heart is to supply new cardiomyocytes differentiated from multipotent cardiovascular progenitor cells or pluripotent stem cells, including embryonic stem cells and induced pluripotent stem cells (iPSC; reviewed extensively elsewhere).<sup>18–21</sup> Cardiomyocytes derived from pluripotent stem cells are a promising source of cells because of recent advances in improving the efficiency of human cardiomyocyte differentiation. However, poor survival, low maturation efficiency, and limited functional integration of engrafted cells are hurdles that must still be overcome. Introduction of adult progenitor cells in animal models has positive effects after myocardial infarction, but the cells do not seem to persist, pointing to a potential paracrine effect.<sup>22,23</sup> Clinical trials are underway to evaluate efficacy, with mixed results to date.<sup>24</sup>


Here, we review a newly emerging strategy for cardiac regeneration that involves reprogramming cells from the vast pool of nonmyocytes in the heart into new cardiomyocytes (Figure 1).<sup>25</sup> This approach builds on recent cellular reprogramming knowledge<sup>26</sup> and leverages >20 years of advances in cardiac developmental biology<sup>20</sup> to induce a cardiomyocyte-like fate in the adult heart. Cardiac reprogramming can be induced by a discrete set of factors but seems to have critical epigenetic blocks in vitro, which are significantly overcome on reprogramming of cells in their native environment in vivo.<sup>25,27–30</sup> We review how advances in cellular reprogramming and developmental biology led to our recent work in direct cardiac reprogramming and consider the future potential of this rapidly evolving approach to cardiac regeneration.



**Figure 1. Approaches to direct cardiac reprogramming.** Several methods for converting nonmyocytes to cardiomyocyte-like cells have been published and are summarized here. In the first example, injection of factors into the noncardiogenic mesoderm of embryonic day (E) 7.0 embryos resulted in ectopic beating cells (E8.5 embryos shown for simplicity). In vitro delivery of various cocktails resulted in primarily partially reprogrammed cells, whereas in vivo delivery yielded more fully reprogrammed cardiomyocyte-like cells. iCMs indicates induced cardiomyocytes.

### Cellular Reprogramming

For decades, the concept in the field of developmental biology had been that cells, once terminally differentiated, were relatively fixed in their cell fate. This dogma was first challenged in the 1960s by the observation that a somatic cell can obtain totipotency through nuclear transfer into an enucleated frog egg.<sup>31</sup> Cloning of Dolly the sheep in 1997 by nuclear transfer revealed that the mammalian egg also had similar ability in reprogramming an adult nucleus to the embryonic state.<sup>32</sup> In the late 1980s and 1990s, further evidence suggested that cell fate conversion can take place by a more direct route. Blau et al<sup>33</sup> demonstrated that fusion of skeletal muscle cells with fibroblasts resulted in heterokaryons that induced conversion of fibroblasts into a skeletal muscle-like phenotype. Studies from the Weintraub laboratory subsequently demonstrated that fibroblasts could be



**Figure 2. Direct conversion of cardiac fibroblasts into cardiomyocyte-like cells in vivo.** Masson trichrome (left) and immunofluorescent staining for  $\alpha$ -Actinin and  $\beta$ -Gal (right) in Gata4, Mef2c, and Tbx5-injected periostin-Cre:R26R-lacZ mouse heart 4 weeks after coronary ligation. Scale bars, 500  $\mu$ m in the left panel, 50  $\mu$ m in the middle and right panels. LV indicates left ventricle; and RV, right ventricle.

converted into skeletal muscle cells *in vitro* with forced expression of the skeletal muscle master regulator gene *MyoD*, which encodes a basic helix-loop-helix (bHLH) domain-containing transcription factor.<sup>34,35</sup> Despite the race to identify individual transcription factors that could function to guide cell fate similar to *MyoD* for other lineages, including cardiomyocytes, the *MyoD* paradigm seemed to be an exception, rather than the rule. Meanwhile, studies in model organisms showed that the forced expression of individual master regulatory transcription factors containing homeobox domains could induce the formation of complex body structures, which is best exemplified by the induction of ectopic eye formation on the legs of *Drosophila* by overexpressing *eyeless*, the fly ortholog of *Pax6*.<sup>36</sup> The concept of cell fate conversion bypassing normal developmental lineage progression emerged but was still largely neglected with respect to the application of such concepts in regenerative medicine.

Takahashi and Yamanaka's<sup>37</sup> milestone publication in 2006 demonstrating the creation of iPSCs ushered in a new era of using cellular reprogramming in regenerative medicine.<sup>38-40</sup> Coexpression of 4 transcription factors (*Oct4*, *Sox2*, *Klf4*, *c-Myc*) was sufficient to reprogram fibroblasts to pluripotent stem cells that had the potential to develop into viable animals.<sup>41,42</sup> Abundant evidence has demonstrated that iPSCs can be differentiated efficiently into multiple cell types that someday could be used for regenerative therapies, disease modeling, and drug discovery. A key lesson from iPSC reprogramming was the recognition that a combinatorial code involving a discrete number of regulatory factors could be sufficient to induce cell fate change.<sup>43,44</sup>

Despite the difficulty in identifying *MyoD*-like factors for direct cellular reprogramming from one terminally differentiated adult somatic cell directly into another without taking a detour back to pluripotency, the iPSC experience raised the possibility of a combination of regulators that could together induce cell fate change or transdifferentiation. This reprogramming strategy was first demonstrated by Zhou et al,<sup>45</sup> who directly converted exocrine pancreatic cells into insulin-producing endocrine cells in the mouse pancreas with the transcription factors *Ngn3*, *Pdx*, and *Mafa*. Similarly, we showed that cardiac and dermal fibroblasts could be reprogrammed into cardiomyocyte-like cells by a combination of 3 transcription factors, *Gata4*, *Mef2c*, and *Tbx5*.<sup>25,27</sup> The direct reprogramming of adult fibroblasts to neuronal-like cells was also achieved *in vitro* with forced expression of a combination of transcription factors<sup>46,47</sup> or microRNAs (miRNAs),<sup>48</sup> and similar observations were made for conversion to the hepatocyte lineage.<sup>49,50</sup> Unlike direct reprogramming with multiple factors, Szabo et al<sup>51</sup> showed that combining the expression of only *Oct4* with administration of select cytokines can reprogram human fibroblasts into hematopoietic progenitors, whereas transient expression of iPSC reprogramming factors, followed by *Jak/Stat* inhibition, resulted in the emergence of cardiomyocytes.<sup>52</sup> The advances in cardiac reprogramming are considered in more detail.

### Discovery of Transcription Factor–Based Direct Cardiac Reprogramming

During the past 20 years, developmental biology studies have revealed complex and intertwined networks of signaling pathways, transcription factors, and miRNAs that regulate the formation and function of the heart.<sup>20,53,54</sup> The networks are self-reinforcing, with

layers of positive and negative feedback loops. Transcription factors often function in common complexes, and human mutations that disrupt their interaction can lead to similar forms of heart malformations as seen with mutations in *GATA4* and *TBX5*.<sup>55</sup> In 2009, Takeuchi and Bruneau<sup>56</sup> demonstrated that overexpression of *Gata4*, *Tbx5*, and the interacting chromatin remodeling protein, *Baf60c*, converts noncardiogenic mesoderm into beating cardiomyocytes in the embryo by a mechanism involving the induction of *Nkx2-5* by *Gata4* and *Baf60c*.

The adult heart has many cell types within the organ that are normally noncardiogenic. The majority of these are cardiac fibroblasts, which comprise >50% of cells in the heart and are derived from an extracardiac structure known as the proepicardial organ.<sup>57</sup> Fibroblasts play an important structural and paracrine role supporting the neighboring myocytes.<sup>58</sup> Upon injury, cardiac fibroblasts are activated and migrate to the site of injury to create scar tissue that replaces dead myocardium. As a result of the abundance of resident cardiac fibroblasts, the ability to reprogram such cells *in situ* would represent a powerful approach for regenerating myocardium.

We leveraged the abundant knowledge of cardiac developmental biology to attempt reprogramming of adult somatic cells into cardiomyocyte-like cells.<sup>25</sup> Despite the desire to ultimately reprogram *in vivo*, we established an *in vitro* assay to discover the minimal cocktail of factors that could convert noncardiomyocytes in the heart into a more cardiomyocyte-like phenotype. We generated transgenic mice containing the EGFP reporter downstream of the cardiomyocyte-specific  $\alpha$ MHC (*Myh6*) promoter<sup>59</sup> and isolated the EGFP-negative nonmyocyte population, which consisted largely of fibroblasts. On retroviral introduction of 14 transcription factors and a pool of miRNAs, neonatal  $\alpha$ MHC-EGFP<sup>-</sup> cells activated the expression of  $\alpha$ MHC (*Myh6*) and the  $\alpha$ MHC-EGFP reporter, which allowed quantitative analysis by fluorescence-activated cell sorting. The miRNAs in this setting were dispensable. Through serial deletion of one transcription factor at a time, we narrowed the required reprogramming factors to the following 3: *Gata4*, *Mef2c*, and *Tbx5* (GMT). The GMT cocktail was sufficient to induce green fluorescent protein (GFP) expression in  $\approx$ 15% to 20% of the cells, which we termed induced cardiomyocytes (iCMs). However, the majority of cells were partially reprogrammed, with 5% of the total infected cell population expressing additional cardiac markers, such as cardiac Troponin T (*cTnT*), and assembling sarcomeric structures; furthermore, only  $\approx$ 0.5% of the  $\alpha$ MHC-EGFP<sup>+</sup>/*cTnT*<sup>+</sup> cells were capable of beating.

Despite the low percentage of fully reprogrammed cardiomyocytes, genome-wide transcriptome studies of the  $\alpha$ MHC-EGFP<sup>+</sup> cells ( $\approx$ 15% of infected cells) showed that the partially reprogrammed population induced a broad cardiac transcriptional program involving hundreds of genes and also silenced the fibroblast transcriptome. Furthermore, the epigenetic status of iCMs was similar to neonatal endogenous cardiomyocytes at loci examined, and the reprogramming event was stable, and did not require ongoing expression of GMT. Notably, the GMT cocktail also reprogrammed tail-tip dermal fibroblasts, albeit with lower efficiency, suggesting that the presence of a cardiac progenitor pool was not necessary for the presence of iCMs.

Interestingly, the more fully reprogrammed iCMs had action potentials that were most similar to adult ventricular myocytes. This observation was in contrast to the relatively immature

electric activity noted in embryonic stem cell–derived or induced pluripotent stem cell–derived cardiomyocytes. Using a Cre-based strategy, we found that iCMs rarely expressed *Mesp1* or *Isl1*, markers of early cardiac progenitors, during the process of cardiac reprogramming.<sup>25</sup> This suggested that the reprogramming event represented a direct conversion from one adult cell type to another rather than traversing through a progenitor stage. The rapidity of initial conversion and the more mature electrophysiology observed in iCMs are consistent with this interpretation.

### In Vivo Direct Cardiac Reprogramming

The relatively poor quality and low efficiency of *in vitro* cardiac reprogramming might be because of the lack of a natural environment for cardiomyocytes on plastic dishes. Because the objective of this strategy was to harness the endogenous cardiac fibroblasts within the organ for regeneration without requiring the use of cell-based therapy, we attempted to deliver GMT *in vivo* after ischemic injury and to convert nonmyocytes to cardiomyocyte-like cells. Classic genetic lineage-tracing studies with periostin-Cre, *Fsp1*-Cre, and  $\alpha$ MHC-MerCreMer were performed to demonstrate that dividing nonmyocytes infected by retroviruses could be converted into iCMs<sup>27</sup> during a period of 4 weeks. Careful studies demonstrated that the newly born myocytes arose from *in vivo* conversion rather than leaky Cre expression or cell–cell fusion. Importantly, intermediate stages of reprogramming to iCMs were identified and characterized at varying time points after retroviral infection, further supporting the idea of a progressive reprogramming process.

*In vivo*–derived iCMs developed many characteristics of endogenous cardiomyocytes. They were binucleate, assembled sarcomeres and had cardiomyocyte-like gene expression by 4 weeks after infection with GMT. Furthermore, single-cell analyses by patch clamp technology revealed that 50% of *in vivo*–derived iCMs closely resembled endogenous cardiomyocytes with beating on electric stimulation and ventricular cardiomyocyte-like action potentials. The markedly improved quality of reprogramming *in vivo* compared with *in vitro* might have been because of signals from the microenvironment, exposure to the extracellular matrix, or influence of mechanical forces while reprogramming. Importantly, we found evidence for electric coupling of the *in vivo* reprogrammed iCMs with endogenous cardiomyocytes and other iCMs. *In vivo* delivery of GMT intramyocardially decreased scar size and attenuated cardiac dysfunction after coronary ligation, as assessed by MRI and echocardiography. As expected, the cardiomyocytes within the scar area of GMT-treated mice represented newly born iCMs as determined by lineage tracing experiments (Figure 2). The beneficial effects of GMT were enhanced with the addition of the proangiogenic and fibroblast-activating factor, thymosin  $\beta$ 4, which independently promotes cardiac repair.<sup>60,61</sup>

In a similar study, Song et al<sup>28</sup> replicated the findings of Ieda et al<sup>25</sup> with GMT and demonstrated that the addition of a bHLH domain–containing transcription factor, *Hand2*, could increase the efficiency of cardiac reprogramming *in vitro*. *Hand2* was initially discovered in a search for a cardiac MyoD-like bHLH protein<sup>62</sup> and is essential for a subset of cardiac progenitors,<sup>63–65</sup> but is not sufficient to induce the cardiac phenotype. Importantly, Song et al<sup>28</sup> also showed direct conversion from nonmyocyte to myocytes *in vivo* by

retrovirally transducing fibroblasts with GMT plus *Hand2* into hearts after coronary ligation. These *in vivo*–induced cardiac-like myocytes were similar to the endogenous cardiomyocytes based on their gene expression, sarcomere structure, and electrophysiological features. Lineage tracing with *Fsp1*-Cre, an inducible *Tcf21*-iCre, and  $\alpha$ MHC-MerCreMer demonstrated that the origin of the iCM-like cells was likely cardiac fibroblasts. This observation was further supported by the evidence that the genetically prelabeled old myocyte pool was diluted by these newly born myocyte-like cells. Along with the emergence of new cardiomyocyte-like cells, improved heart function and reduced scar size were observed.

A third successful example of *in vivo* cardiac reprogramming was reported by Inagawa et al.<sup>30</sup> The authors used the GMT cocktail and retroviral delivery method. Different from Qian et al<sup>27</sup> and Song et al,<sup>28</sup> immunosuppressed mice were used in an attempt to promote the survival of viral-transduced cells. Reprogramming with these nude mice resulted in more iCMs with well-defined sarcomere structure. Furthermore, to improve the transduction of GMT *in vivo*, Inagawa et al<sup>30</sup> generated a polycistronic retrovirus expressing GMT at near-equimolar levels from the same promoter using self-cleaving 2A peptides that resulted in a better reprogramming efficiency. Although the global outcome of GMT introduction in this system was not clear, use of a 2A polycistronic vector and immunosuppressed mice further refined the *in vivo* reprogramming technology.

These 3 studies provide compelling evidence that the abundant nonmyocyte pool in the heart, largely composed of fibroblasts, can be transdifferentiated into new cardiomyocyte-like cells *in vivo* after injury, resulting in regeneration of myocardium and improved cardiac function. The functional improvement can be partially explained by new myocytes that increase force generation, but a fundamental alteration in the nature of scar-producing fibroblasts could also account for part of the reduction in scar size and improvement in cardiac output.

### Alternative Strategies for Cardiac Reprogramming

The strategy involving serial deletion of 1 factor at a time can yield potential combinations of factors for reprogramming but, because of the complex interactions between networks, may not necessarily yield the optimal combination. To address this, Protze et al<sup>66</sup> screened 120 triplet combinations of 10 important developmental cardiac transcription factors expressed by lentiviruses in mouse embryonic fibroblasts for their ability to induce a myocyte-like phenotype. Rather than measuring the activation of a single reporter, they assayed expression of a panel of cardiac genes by quantitative polymerase chain reaction. In this screen, myocardin, *Mef2c*, and *Tbx5* comprised the optimal combination to convert fibroblasts into myocyte-like cells, determined by gene expression, sarcomere formation, and ion channel activity, suggesting multiple combinations may exist for efficient cardiac reprogramming. Protze et al<sup>66</sup> also performed a time-course experiment and showed a progressive process involved in direct cardiac reprogramming in which a more complete cardiac phenotype arose over time.

miRNAs have important roles in cardiomyocyte decisions and are often regulated by the major cardiac transcription factors and, in turn, titrate the dosage of the key transcriptional networks.<sup>54,67</sup> In particular, miR-1, the most abundant cardiac miRNA, promotes

muscle gene expression and regulates many aspects of cardiac biology.<sup>68–71</sup> Recently, Jayawardena et al<sup>29</sup> showed that a combination of 4 muscle-specific miRNAs (miR-1, miR-133, miR-208, miR-499) was sufficient to transdifferentiate mouse fibroblasts into cardiomyocyte-like cells *in vitro* and *in vivo*. Remarkably, in the presence of a Janus kinase (JAK)-1 inhibitor, miR-1 alone was sufficient to reprogram the cardiac fibroblasts.<sup>29</sup>

Although many groups have now successfully reprogrammed fibroblasts to cardiomyocyte-like cells *in vitro*, others have struggled,<sup>72</sup> highlighting the challenges to easily mastering this technique. The use of fresh, nonsenescent fibroblasts, high titers of viruses expressing the reprogramming factors, and careful culture conditions are among the variables involved in achieving successful reprogramming.<sup>73,74</sup> Future work on standardizing the conditions and improving the conversion efficiency *in vitro* will be necessary to make this a routine procedure in many other laboratories.

### Reprogramming Human Fibroblasts Toward the Cardiomyocyte Fate

In order to advance cardiac reprogramming technology, it will be important to translate the knowledge gained from studies using the mouse system into human cells. Several recent studies reported the failure of GMT or GMHT to convert fibroblasts to cardiomyocyte-like cells, but each described overlapping but distinct combinations of factors that could push human fibroblasts into a more cardiomyocyte-like state.<sup>75–77</sup> Nam et al<sup>75</sup> reported that a combination of 4 transcription factors (Gata-4, Hand2, Tbx-5, and Myocd) and 2 miRNAs (miR-1 and miR-133) could reprogram up to 20% of human fibroblasts into cTnT-expressing cells. Upregulation of cardiac genes and downregulation of fibroblast genes was observed, and some cells developed calcium transients with observation of rare beating cells. However, many cardiac genes failed to be upregulated, suggesting the switch from a fibroblast-like phenotype to cardiomyocyte-like phenotype is partial. Similarly, Fu et al<sup>76</sup> found that addition of MESP1 and myocardin to GMT, along with a nuclear hormone receptor, ESRRG, could induce gene expression shifts in human fibroblasts to a similar extent as GMT in mouse cells *in vitro*. Approximately 20% of these cells developed calcium transients, and some had action potentials similar to human pluripotent stem cell-derived cardiomyocytes. Wada et al<sup>77</sup> also found that addition of MESP1 and Myocardin to GMT could reprogram human fibroblasts into cells that had many properties of human cardiomyocytes. Thus, while human cells appear more resistant to conversion, various combinations of the core transcriptional machinery are able to shift cells considerably toward the cardiomyocyte state, although further improvements may be necessary. Whether the current combinations of reprogramming factors result in more complete reprogramming *in vivo*, similar to the experience in mouse cells, await studies in large animals, including non-human primates.

### Challenges and Future Directions

Despite the excitement and potential of direct cardiac reprogramming technology, several challenges remain, as would be expected for any new technology. These challenges include developing a mechanistic understanding of the reprogramming process, improving efficiency in mouse and human cells, enhancement or replacement of genes with drug-like molecules,

and performing *in vivo* efficacy and safety in large animals. Such efforts are underway in many labs, and the coming years should rapidly yield advancements in this nascent technology for regenerative medicine.

The low conversion rate of fibroblasts to fully reprogrammed iCMs *in vitro* is a major challenge for deciphering the mechanism of reprogramming. The relative inefficiency *in vitro* is not surprising, considering the iPSC reprogramming efficiency rate (0.01%–0.1%), which is likely because of major epigenetic barriers that cells cannot easily overcome.<sup>78</sup> Unlike iPSCs, reprogrammed cardiomyocytes rapidly exit the cell cycle, making efficiency a much bigger concern. Understanding how progressive epigenetic and transcriptional changes occur temporally during direct cardiac reprogramming is an important first step toward overcoming these hurdles. As highlighted from 2 recent genome-wide epigenetic studies of embryonic stem cell (ESC)-derived cardiomyocytes,<sup>79,80</sup> temporal alterations in chromatin structure patterns lead to activation of key genes associated with heart development and function at distinctive differentiation stages. Such mechanisms might be informative for cardiac reprogramming. Second, methods aiming to improve delivery efficiency, such as the use of a polycistronic vector<sup>30</sup> to control for homogenous reprogramming gene expression, nanoparticle techniques for specific and efficient targeting, immunosuppression to promote survival of transduced cells,<sup>30</sup> the administration of thymosin  $\beta$ 4 to mobilize and activate fibroblasts,<sup>60</sup> and use of VEGF to promote angiogenesis,<sup>81</sup> may increase the final number of reprogrammed iCMs. Third, techniques that enhance the maturation of cardiomyocytes should be used to promote the progression of iCMs from immature to mature stages. These include supplying the cells with cytokines at defined stages, altering the expression pattern of certain ion channel regulators, exposing cells to mechanical forces, and providing extracellular matrix and endothelial cells. Finally, small-molecule and secreted factor screens geared toward increasing iCM numbers and quality should be performed to identify externally administered factors that promote the reprogramming process.

Approaches to improve efficiency *in vitro* will aid in mechanistic understanding of the reprogramming process and may ultimately allow use of direct reprogramming to model human disease. Although current technology does not generate sufficient numbers of fully reprogrammed iCMs for disease modeling studies, improvements in efficiency will be valuable because iCMs seem to achieve electric maturation that is more similar to ventricular myocytes, particularly when reprogrammed *in vivo*, and this has been difficult to achieve with ESC-derived or iPSC-derived cardiomyocytes. Generation of sufficient numbers of fully reprogrammed cells *in vitro* would also be valuable for drug toxicity studies and drug screening.

In contrast, current iCM technology is quite efficient for *in vivo* reprogramming. Thus, harnessing the vast endogenous pool of noncardiomyocytes seems like a viable approach to regenerate heart muscle without cell-based therapy. Although this would require a gene therapy approach ideally using non-integration vectors such as adeno-associated viral vectors, a reasonable regulatory path exists for virally mediated gene delivery, with scores of Food and Drug Administration–approved trials underway and the recent approval of a gene therapy drug

for lipoprotein lipase deficiency in Europe. Future identification of small molecules or secreted proteins that could replace each transcription factor, as has been performed for iPSC reprogramming,<sup>82,83</sup> may allow an alternative to gene therapy. It is possible that in vivo reprogramming of cells to regenerate damaged tissue will serve as a new paradigm for many human diseases, and the lessons learned in the cardiac area will be applicable to strategies to realize this dream in other tissues. Although many challenges lie ahead in advancing this nascent technology, the opportunities and the potential benefits are significant, and we are confident that the field will continue to push this technology further in the years ahead.

### Acknowledgments

We thank the members of the Srivastava laboratory for valuable discussion and contributions, B. Taylor for assistance with preparation of the article, and G. Howard for editorial assistance.

### Sources of Funding

D. Srivastava is supported by grants from National Heart, Lung, and Blood Institute/National Institutes of Health, the California Institute for Regenerative Medicine, the Younger Family Foundation, the Roddenberry Foundation, and the L.K. Whittier Foundation.

### Disclosures

D. Srivastava serves on the Scientific Advisory Board of iPierian, Inc, Berkeley Lights, Inc, and RegeneRx Biopharmaceuticals, Inc. The other author reports no conflicts.

### References

- Olivetti G, Melissari M, Capasso JM, Anversa P. Cardiomyopathy of the aging human heart. Myocyte loss and reactive cellular hypertrophy. *Circ Res*. 1991;68:1560–1568.
- Poss KD, Wilson LG, Keating MT. Heart regeneration in zebrafish. *Science*. 2002;298:2188–2190.
- Soonpaa MH, Field LJ. Assessment of cardiomyocyte DNA synthesis in normal and injured adult mouse hearts. *Am J Physiol*. 1997;272:H220–H226.
- Soonpaa MH, Field LJ. Survey of studies examining mammalian cardiomyocyte DNA synthesis. *Circ Res*. 1998;83:15–26.
- Senyo SE, Steinhauser ML, Pizzimenti CL, Yang VK, Cai L, Wang M, Wu TD, Guerin-Kern JL, Lechene CP, Lee RT. Mammalian heart renewal by pre-existing cardiomyocytes. *Nature*. 2013;493:433–436.
- Bergmann O, Bhardwaj RD, Bernard S, Zdunek S, Barnabé-Heider F, Walsh S, Zupcic H, Alkass K, Buchholz BA, Druid H, Jovinge S, Frisén J. Evidence for cardiomyocyte renewal in humans. *Science*. 2009;324:98–102.
- Quaini F, Urbanek K, Graiani G, Lagrasta C, Maestri R, Monica M, Boni A, Ferraro F, Delsignore R, Tasca G, Leri A, Kajstura J, Quaini E, Anversa P. The regenerative potential of the human heart. *Int J Cardiol*. 2004;95(Suppl 1):S26–S28.
- Hsieh PC, Segers VF, Davis ME, MacGillivray C, Gannon J, Molkenin JD, Robbins J, Lee RT. Evidence from a genetic fate-mapping study that stem cells refresh adult mammalian cardiomyocytes after injury. *Nat Med*. 2007;13:970–974.
- Porrello ER, Mahmoud AI, Simpson E, Hill JA, Richardson JA, Olson EN, Sadek HA. Transient regenerative potential of the neonatal mouse heart. *Science*. 2011;331:1078–1080.
- Kikuchi K, Holdway JE, Werdich AA, Anderson RM, Fang Y, Egnaczyk GF, Evans T, Macrae CA, Stainier DY, Poss KD. Primary contribution to zebrafish heart regeneration by *gata4*(+) cardiomyocytes. *Nature*. 2010;464:601–605.
- Jopling C, Sleep E, Raya M, Martí M, Raya A, Izpisua Belmonte JC. Zebrafish heart regeneration occurs by cardiomyocyte dedifferentiation and proliferation. *Nature*. 2010;464:606–609.
- Hassink RJ, Pasumarthi KB, Nakajima H, Rubart M, Soonpaa MH, de la Rivière AB, Doevendans PA, Field LJ. Cardiomyocyte cell cycle activation improves cardiac function after myocardial infarction. *Cardiovasc Res*. 2008;78:18–25.
- Soonpaa MH, Koh GY, Pajak L, Jing S, Wang H, Franklin MT, Kim KK, Field LJ. Cyclin D1 overexpression promotes cardiomyocyte DNA synthesis and multinucleation in transgenic mice. *J Clin Invest*. 1997;99:2644–2654.
- Kühn B, del Monte F, Hajjar RJ, Chang YS, Lebeche D, Arab S, Keating MT. Periostin induces proliferation of differentiated cardiomyocytes and promotes cardiac repair. *Nat Med*. 2007;13:962–969.
- Engel FB, Hsieh PC, Lee RT, Keating MT. FGF1/p38 MAP kinase inhibitor therapy induces cardiomyocyte mitosis, reduces scarring, and rescues function after myocardial infarction. *Proc Natl Acad Sci U S A*. 2006;103:15546–15551.
- Bersell K, Arab S, Haring B, Kühn B. Neuregulin1/ErbB4 signaling induces cardiomyocyte proliferation and repair of heart injury. *Cell*. 2009;138:257–270.
- Eulalio A, Mano M, Dal Ferro M, Zentilin L, Sinagra G, Zacchigna S, Giacca M. Functional screening identifies miRNAs inducing cardiac regeneration. *Nature*. 2012;492:376–381.
- Murry CE, Keller G. Differentiation of embryonic stem cells to clinically relevant populations: lessons from embryonic development. *Cell*. 2008;132:661–680.
- Passier R, van Laake LW, Mummery CL. Stem-cell-based therapy and lessons from the heart. *Nature*. 2008;453:322–329.
- Srivastava D, Ivey KN. Potential of stem-cell-based therapies for heart disease. *Nature*. 2006;441:1097–1099.
- Burridge PW, Keller G, Gold JD, Wu JC. Production of de novo cardiomyocytes: human pluripotent stem cell differentiation and direct reprogramming. *Cell Stem Cell*. 2012;10:16–28.
- Mummery CL, Davis RP, Krieger JE. Challenges in using stem cells for cardiac repair. *Sci Transl Med*. 2010;2:27ps17.
- Lafamme MA, Murry CE. Heart regeneration. *Nature*. 2011;473:326–335.
- Mathur A, Martin JF. Stem cells and repair of the heart. *Lancet*. 2004;364:183–192.
- Ieda M, Fu JD, Delgado-Olguin P, Vedantham V, Hayashi Y, Bruneau BG, Srivastava D. Direct reprogramming of fibroblasts into functional cardiomyocytes by defined factors. *Cell*. 2010;142:375–386.
- Yamanaka S. Induced pluripotent stem cells: past, present, and future. *Cell Stem Cell*. 2012;10:678–684.
- Qian L, Huang Y, Spencer CL, Foley A, Vedantham V, Liu L, Conway SJ, Fu JD, Srivastava D. In vivo reprogramming of murine cardiac fibroblasts into induced cardiomyocytes. *Nature*. 2012;485:593–598.
- Song K, Nam YJ, Luo X, Qi X, Tan W, Huang GN, Acharya A, Smith CL, Tallquist MD, Neilson EG, Hill JA, Bassel-Duby R, Olson EN. Heart repair by reprogramming non-myocytes with cardiac transcription factors. *Nature*. 2012;485:599–604.
- Jayawardena TM, Egemazarov B, Finch EA, Zhang L, Payne JA, Pandya K, Zhang Z, Rosenberg P, Mirotsov M, Dzau VJ. MicroRNA-mediated in vitro and in vivo direct reprogramming of cardiac fibroblasts to cardiomyocytes. *Circ Res*. 2012;110:1465–1473.
- Inagawa K, Miyamoto K, Yamakawa H, Muraoka N, Sadahiro T, Umei T, Wada R, Katsumata Y, Kaneda R, Nakade K, Kurihara C, Obata Y, Miyake K, Fukuda K, Ieda M. Induction of cardiomyocyte-like cells in infarct hearts by gene transfer of *Gata4*, *Mef2c*, and *Tbx5*. *Circ Res*. 2012;111:1147–1156.
- Gurdon JB. The developmental capacity of nuclei taken from intestinal epithelium cells of feeding tadpoles. *J Embryol Exp Morphol*. 1962;10:622–640.
- Gurdon JB. From nuclear transfer to nuclear reprogramming: the reversal of cell differentiation. *Annu Rev Cell Dev Biol*. 2006;22:1–22.
- Blau HM, Pavlath GK, Hardeman EC, Chiu CP, Silberstein L, Webster SG, Miller SC, Webster C. Plasticity of the differentiated state. *Science*. 1985;230:758–766.
- Lassar AB, Paterson BM, Weintraub H. Transfection of a DNA locus that mediates the conversion of 10T1/2 fibroblasts to myoblasts. *Cell*. 1986;47:649–656.
- Weintraub H, Tapscott SJ, Davis RL, Thayer MJ, Adam MA, Lassar AB, Miller AD. Activation of muscle-specific genes in pigment, nerve, fat, liver, and fibroblast cell lines by forced expression of MyoD. *Proc Natl Acad Sci U S A*. 1989;86:5434–5438.
- Halder G, Callaerts P, Gehring WJ. Induction of ectopic eyes by targeted expression of the *eyeless* gene in *Drosophila*. *Science*. 1995;267:1788–1792.
- Takahashi K, Yamanaka S. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell*. 2006;126:663–676.
- Takahashi K, Tanabe K, Ohnuki M, Narita M, Ichisaka T, Tomoda K, Yamanaka S. Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell*. 2007;131:861–872.

39. Yu J, Vodyanik MA, Smuga-Otto K, Antosiewicz-Bourget J, Frane JL, Tian S, Nie J, Jonsdottir GA, Ruotti V, Stewart R, Slukvin II, Thomson JA. Induced pluripotent stem cell lines derived from human somatic cells. *Science*. 2007;318:1917–1920.
40. Park IH, Zhao R, West JA, Yabuuchi A, Huo H, Ince TA, Lerou PH, Lensch MW, Daley GQ. Reprogramming of human somatic cells to pluripotency with defined factors. *Nature*. 2008;451:141–146.
41. Zhao XY, Li W, Lv Z, Liu L, Tong M, Hai T, Hao J, Guo CL, Ma QW, Wang L, Zeng F, Zhou Q. iPS cells produce viable mice through tetraploid complementation. *Nature*. 2009;461:86–90.
42. Boland MJ, Hazen JL, Nazor KL, Rodriguez AR, Gifford W, Martin G, Kupriyanov S, Baldwin KK. Adult mice generated from induced pluripotent stem cells. *Nature*. 2009;461:91–94.
43. Sridharan R, Tchieu J, Mason MJ, Yachechko R, Kuoy E, Horvath S, Zhou Q, Plath K. Role of the murine reprogramming factors in the induction of pluripotency. *Cell*. 2009;136:364–377.
44. Buganim Y, Faddah DA, Cheng AW, Itskovich E, Markoulaki S, Ganz K, Klemm SL, van Oudenaarden A, Jaenisch R. Single-cell expression analyses during cellular reprogramming reveal an early stochastic and a late hierarchic phase. *Cell*. 2012;150:1209–1222.
45. Zhou Q, Brown J, Kanarek A, Rajagopal J, Melton DA. In vivo reprogramming of adult pancreatic exocrine cells to beta-cells. *Nature*. 2008;455:627–632.
46. Pang ZP, Yang N, Vierbuchen T, Ostermeier A, Fuentes DR, Yang TQ, Citri A, Sebastiano V, Marro S, Südhof TC, Wernig M. Induction of human neuronal cells by defined transcription factors. *Nature*. 2011;476:220–223.
47. Vierbuchen T, Ostermeier A, Pang ZP, Kokubu Y, Südhof TC, Wernig M. Direct conversion of fibroblasts to functional neurons by defined factors. *Nature*. 2010;463:1035–1041.
48. Yoo AS, Sun AX, Li L, Shcheglovitov A, Portmann T, Li Y, Lee-Messer C, Dolmetsch RE, Tsien RW, Crabtree GR. MicroRNA-mediated conversion of human fibroblasts to neurons. *Nature*. 2011;476:228–231.
49. Huang P, He Z, Ji S, Sun H, Xiang D, Liu C, Hu Y, Wang X, Hui L. Induction of functional hepatocyte-like cells from mouse fibroblasts by defined factors. *Nature*. 2011;475:386–389.
50. Sekiya S, Suzuki A. Direct conversion of mouse fibroblasts to hepatocyte-like cells by defined factors. *Nature*. 2011;475:390–393.
51. Szabo E, Rampalli S, Risueño RM, Schnerch A, Mitchell R, Fiebig-Comyn A, Levadoux-Martin M, Bhatia M. Direct conversion of human fibroblasts to multilineage blood progenitors. *Nature*. 2010;468:521–526.
52. Efe JA, Hilcove S, Kim J, Zhou H, Ouyang K, Wang G, Chen J, Ding S. Conversion of mouse fibroblasts into cardiomyocytes using a direct reprogramming strategy. *Nat Cell Biol*. 2011;13:215–222.
53. Olson EN. Gene regulatory networks in the evolution and development of the heart. *Science*. 2006;313:1922–1927.
54. Cordes KR, Srivastava D. MicroRNA regulation of cardiovascular development. *Circ Res*. 2009;104:724–732.
55. Garg V, Kathiriyai IS, Barnes R, Schluterman MK, King IN, Butler CA, Rothrock CR, Eapen RS, Hirayama-Yamada K, Joo K, Matsuoka R, Cohen JC, Srivastava D. GATA4 mutations cause human congenital heart defects and reveal an interaction with TBX5. *Nature*. 2003;424:443–447.
56. Takeuchi JK, Bruneau BG. Directed transdifferentiation of mouse mesoderm to heart tissue by defined factors. *Nature*. 2009;459:708–711.
57. Snider P, Hinton RB, Moreno-Rodriguez RA, Wang J, Rogers R, Lindsley A, Li F, Ingram DA, Menick D, Field L, Firulli AB, Molkentin JD, Markwald R, Conway SJ. Periostin is required for maturation and extracellular matrix stabilization of noncardiomyocyte lineages of the heart. *Circ Res*. 2008;102:752–760.
58. Ieda M, Tsuchihashi T, Ivey KN, Ross RS, Hong TT, Shaw RM, Srivastava D. Cardiac fibroblasts regulate myocardial proliferation through beta1 integrin signaling. *Dev Cell*. 2009;16:233–244.
59. Gulick J, Subramaniam A, Neumann J, Robbins J. Isolation and characterization of the mouse cardiac myosin heavy chain genes. *J Biol Chem*. 1991;266:9180–9185.
60. Smart N, Risebro CA, Melville AA, Moses K, Schwartz RJ, Chien KR, Riley PR. Thymosin beta4 induces adult epicardial progenitor mobilization and neovascularization. *Nature*. 2007;445:177–182.
61. Bock-Marquette I, Saxena A, White MD, Dimaio JM, Srivastava D. Thymosin beta4 activates integrin-linked kinase and promotes cardiac cell migration, survival and cardiac repair. *Nature*. 2004;432:466–472.
62. Srivastava D, Cserjesi P, Olson EN. A subclass of bHLH proteins required for cardiac morphogenesis. *Science*. 1995;270:1995–1999.
63. Srivastava D, Thomas T, Lin Q, Kirby ML, Brown D, Olson EN. Regulation of cardiac mesodermal and neural crest development by the bHLH transcription factor, dHAND. *Nat Genet*. 1997;16:154–160.
64. Yamagishi H, Yamagishi C, Nakagawa O, Harvey RP, Olson EN, Srivastava D. The combinatorial activities of Nkx2.5 and dHAND are essential for cardiac ventricle formation. *Dev Biol*. 2001;239:190–203.
65. Tsuchihashi T, Maeda J, Shin CH, Ivey KN, Black BL, Olson EN, Yamagishi H, Srivastava D. Hand2 function in second heart field progenitors is essential for cardiogenesis. *Dev Biol*. 2011;351:62–69.
66. Protze S, Khattak S, Poulet C, Lindemann D, Tanaka EM, Ravens U. A new approach to transcription factor screening for reprogramming of fibroblasts to cardiomyocyte-like cells. *J Mol Cell Cardiol*. 2012;53:323–332.
67. van Rooij E, Olson EN. MicroRNAs: powerful new regulators of heart disease and provocative therapeutic targets. *J Clin Invest*. 2007;117:2369–2376.
68. Lim LP, Lau NC, Garrett-Engle P, Grimson A, Schelter JM, Castle J, Bartel DP, Linsley PS, Johnson JM. Microarray analysis shows that some microRNAs downregulate large numbers of target mRNAs. *Nature*. 2005;433:769–773.
69. Zhao Y, Samal E, Srivastava D. Serum response factor regulates a muscle-specific microRNA that targets Hand2 during cardiogenesis. *Nature*. 2005;436:214–220.
70. Zhao Y, Ransom JF, Li A, Vedantham V, von Drehle M, Muth AN, Tsuchihashi T, McManus MT, Schwartz RJ, Srivastava D. Dysregulation of cardiogenesis, cardiac conduction, and cell cycle in mice lacking miRNA-1-2. *Cell*. 2007;129:303–317.
71. Ivey KN, Muth A, Arnold J, King FW, Yeh RF, Fish JE, Hsiao EC, Schwartz RJ, Conklin BR, Bernstein HS, Srivastava D. MicroRNA regulation of cell lineages in mouse and human embryonic stem cells. *Cell Stem Cell*. 2008;2:219–229.
72. Chen JX, Krane M, Deutsch MA, Wang L, Rav-Acha M, Gregoire S, Engels MC, Rajarajan K, Karra R, Abel ED, Wu JC, Milan D, Wu SM. Inefficient reprogramming of fibroblasts into cardiomyocytes using Gata4, Mef2c, and Tbx5. *Circ Res*. 2012;111:50–55.
73. Srivastava D, Ieda M. Critical factors for cardiac reprogramming. *Circ Res*. 2012;111:5–8.
74. Qian L, Berry EC, Fu JD, Ieda M, Srivastava D. Reprogramming of mouse fibroblasts to cardiomyocyte-like cells in vitro. *Nat Protocol*. 2013;8:1204–1215.
75. Nam YJ, Song K, Luo X, Daniel E, Lambeth K, West K, Hill JA, DiMaio JM, Baker LA, Bassel-Duby R, Olson EN. Reprogramming of human fibroblasts toward a cardiac fate. *Proc Natl Acad Sci USA*. 2013;110:5588–5593.
76. Fu J, Stone NR, Liu L, Spencer CI, Qian L, Hayashi Y, Delgado-Olguin P, Ding S, Bruneau BG, Srivastava D. Direct reprogramming of human fibroblasts toward the cardiomyocyte lineage. *Stem Cell Reports* (in press).
77. Wada R, Muraoka N, Inagawa K, Yamakawa H, Miyamoto K, Sadahiro T, Umei T, Kaneda R, Suzuki T, Kamiya K, Tohyama S, Yuasa S, Kokaji K, Aeba R, Yozu R, Yamagishi H, Kitamura T, Fukuda K, Ieda M. Induction of human cardiomyocyte-like cells from fibroblasts by defined factors. *Proc Natl Acad Sci USA*. 2013;110:12667–12672.
78. Plath K, Lowry WE. Progress in understanding reprogramming to the induced pluripotent state. *Nat Rev Genet*. 2011;12:253–265.
79. Wamstad JA, Alexander JM, Truty RM, et al. Dynamic and coordinated epigenetic regulation of developmental transitions in the cardiac lineage. *Cell*. 2012;151:206–220.
80. Paige SL, Thomas S, Stoick-Cooper CL, Wang H, Maves L, Sandstrom R, Pabon L, Reinecke H, Pratt G, Keller G, Moon RT, Stamatoyannopoulos J, Murry CE. A temporal chromatin signature in human embryonic stem cells identifies regulators of cardiac development. *Cell*. 2012;151:221–232.
81. Mathison M, Gersch RP, Nasser A, Lilo S, Korman M, Fourman M, Hackett N, Shroyer K, Yang J, Ma Y, Crystal RG, Rosengart TK. In vivo cardiac cellular reprogramming efficacy is enhanced by angiogenic preconditioning of the infarcted myocardium with vascular endothelial growth factor. *J Am Heart Assoc*. 2012;1:e005652.
82. Zhou H, Wu S, Joo JY, Zhu S, Han DW, Lin T, Trauger S, Bien G, Yao S, Zhu Y, Siuzdak G, Schöler HR, Duan L, Ding S. Generation of induced pluripotent stem cells using recombinant proteins. *Cell Stem Cell*. 2009;4:381–384.
83. Hou P, Li Y, Zhang X, Liu C, Guan J, Li H, Zhao T, Ye J, Yang W, Liu K, Ge J, Xu J, Zhang Q, Zhao Y, Deng H. Pluripotent stem cells induced from mouse somatic cells by small-molecule compounds. *Science*. 2013;341:651–654.